



TITLE:

Construction of solutions  $f'(x) = af(\lambda x), \lambda > 1$ . (Banach spaces, function spaces, inequalities and their applications)

AUTHOR(S):

Yoneda, Tsuyoshi

---

CITATION:

Yoneda, Tsuyoshi. Construction of solutions  $f'(x) = af(\lambda x), \lambda > 1$ . (Banach spaces, function spaces, inequalities and their applications). 数理解析研究所講究録 2007, 1570: 1-7

ISSUE DATE:

2007-10

URL:

<http://hdl.handle.net/2433/81278>

RIGHT:

# Construction of solutions $f'(x) = af(\lambda x)$ , $\lambda > 1$ .

東大数理 米田 剛 (Tsuyoshi Yoneda)

Graduate School of Mathematical Sciences

The University of Tokyo

## 1. MAIN RESULT

We denote by  $C^\infty(\mathbb{R})$  the set of all infinitely differentiable functions on  $\mathbb{R}$ . Let

$$C_0^\infty(\mathbb{R}) = \left\{ f \in C^\infty(\mathbb{R}) : \frac{d^\ell f}{dx^\ell}(x) \rightarrow 0 \text{ as } |x| \rightarrow \infty, \ell = 0, 1, 2, \dots \right\},$$

$$C_{\text{comp}}^\infty(\mathbb{R}) = \{f \in C^\infty(\mathbb{R}) : f \text{ has a compact support}\}.$$

We denote by  $\chi_{[r,s]}$  the characteristic function of the interval  $[r, s]$ . Let  $X$  be the set of all step functions

$$(1.1) \quad p = \sum_{j=1}^m c_j \chi_{[r_{j-1}, r_j]},$$

such that  $m = 1, 2, \dots$  and

$$c_j \in \mathbb{R} \quad (j = 1, 2, \dots, m), \quad c_1 = 1,$$

$$r_0 = 0, \quad 0 < r_1 < r_2 < \dots < r_m < \infty,$$

$$\sum_{j=1}^m c_j(r_j - r_{j-1}) = 1.$$

We note that  $\int_{\mathbb{R}} p(x) dx = 1$  for  $p \in X$ .

For  $\lambda > 1$  and  $p \in X$ , we define an operator  $T = T_{\lambda,p} : L^1(\mathbb{R}) \rightarrow L^1(\mathbb{R})$  (by [17] (p. 149, Remark 2) and [2], we can say that  $T_{\lambda,p} : B_{1,\infty}^s(\mathbb{R}) \rightarrow B_{1,\infty}^{s+1}(\mathbb{R})$  for  $s \in \mathbb{R}$ , since  $\lambda p(\lambda \cdot) \in B_{1,\infty}^1(\mathbb{R})$ ) as follows:

$$(1.2) \quad Tf(x) = T_{\lambda,p}f(x) = \lambda(p * f)(\lambda x).$$

We note that  $Tf \in L^1(\mathbb{R}) \cap C(\mathbb{R})$ , since  $f \in L^1(\mathbb{R})$  and  $p \in L_{\text{comp}}^\infty(\mathbb{R})$ .

If  $f \in B_{1,\infty}^s(\mathbb{R})$  we have  $\hat{f}(1 + |\cdot|^2)^{s/2} \in L^\infty(\mathbb{R})$  (see [15]). Therefore we have  $T^k f \in L^2(\mathbb{R}) \cap C(\mathbb{R})$  for  $k > -s - 1$  if  $f \in B_{1,\infty}^s(\mathbb{R})$ .

**Theorem 1.1.** For  $\lambda > 1$  and  $p \in X$ , let  $T$  be the operator defined by (1.2). Then there exists a function  $u = u_{\lambda,p} \in C_{\text{comp}}^\infty(\mathbb{R})$  such that, for all  $f \in L^1(\mathbb{R})$  (or for all  $f \in \cup_{s \in \mathbb{R}} B_{1,\infty}^s(\mathbb{R})$ , which function set is strictly bigger than  $L^1(\mathbb{R})$ ),

$$\lim_{\ell \rightarrow \infty} T^\ell f(x) = c_f u(x) \quad \text{uniformly with respect to } x \in \mathbb{R},$$

$$\text{where } c_f = \hat{f}(0).$$

Moreover,

- (1)  $Tu = u$ ;
- (2)  $\int_{\mathbb{R}} u(x) dx = 1$ ;
- (3)  $\frac{d^k}{dx^k} u(0) = 0$  for all  $k = 0, 1, 2, \dots$ ;
- (4) if  $p \geq 0$ , then  $u \geq 0$ ; and,
- (5) if  $\text{supp } p \subset [0, r]$ , then  $\text{supp } u \subset [0, r/(\lambda - 1)]$ .

In the above statement  $\text{supp } f$  denotes the support of  $f$ .

**Remark 1.1.** If  $p_1 \neq p_2$ , then  $u_{\lambda,p_1} \neq u_{\lambda,p_2}$ .

## 2. APPLICATION

To construct solutions for the following equation,

$$(2.1) \quad \begin{cases} f'(x) = \lambda^2 f(\lambda x), & x \in \mathbb{R}, \\ f(0) = 0, \end{cases}$$

we use  $u = u_{\lambda,p}$  in Theorem 1.1 with  $\lambda > 1$  and  $p = \sum_{j=1}^m c_j \chi_{[r_{j-1}, r_j]} \in X$ , and we define a sequence of functions  $\{v_\ell\}$  inductively as follows:

$$\begin{cases} v_0(x) = u(x) + \sum_{j=1}^m \tilde{c}_j u(x - r_j), \\ v_{\ell+1}(x) = v_\ell(x) + \sum_{j=1}^m \tilde{c}_j v_\ell(x - \lambda^{\ell+1} r_j), \quad \ell = 0, 1, 2, \dots, \end{cases}$$

where  $\tilde{c}_j = -(c_j - c_{j+1})$ ,  $j = 1, 2, \dots, m-1$ ,  $\tilde{c}_m = -c_m$ .

Then  $v_\ell \in C_{\text{comp}}^\infty(\mathbb{R})$ ,  $\ell = 0, 1, 2, \dots$ , and

$$v_\ell(x) = v_{\ell+1}(x), \quad x \in (-\infty, \lambda^{\ell+1} r_1], \quad \ell = 0, 1, 2, \dots$$

Therefore we get a function  $f = f_{\lambda,p} \in C^\infty(\mathbb{R})$  such that

$$f(x) = v_\ell(x), \quad x \in (-\infty, \lambda^{\ell+1} r_1], \quad \ell = 0, 1, 2, \dots$$

Since  $f(x) = u(x)$  near  $x = 0$ ,  $\frac{d^k}{dx^k} f(0) = 0$  for all  $k = 0, 1, 2, \dots$ .

**Theorem 2.1.** *Let  $\lambda > 1$ . For every  $p \in X$ , the function  $f = f_{\lambda,p}$  given by the above method is a solution for (2.1).*

*Remark 2.1.* By the property (5) in Theorem 1.1 and the definition of  $v_\ell$ ,

$$\begin{aligned}\operatorname{supp} u &\subset [0, r_m/(\lambda - 1)], \\ \operatorname{supp}(v_0 - u) &\subset [r_1, r_m\lambda/(\lambda - 1)], \\ \operatorname{supp} v_\ell &\subset [0, r_m\lambda^{\ell+1}/(\lambda - 1)], \\ \operatorname{supp}(v_{\ell+1} - v_\ell) &\subset [r_1\lambda^{\ell+1}, r_m\lambda^{\ell+2}/(\lambda - 1)].\end{aligned}$$

Therefore, if  $r_m/(\lambda - 1) < r_1$ , then  $f(x) = 0$  on  $[r_m\lambda^\ell/(\lambda - 1), r_1\lambda^\ell]$ ,  $\ell = 0, 1, 2, \dots$ , and  $f$  is bounded (See the graph (S12)). Let

$$p_i = \sum_{j=1}^{m_i} c_{i,j} \chi_{[r_{i,j-1}, r_{i,j}]} \in X, \quad i = 1, 2.$$

If  $p_1 \neq p_2$  and there exists  $R > 0$  such that  $r_{i,m_i}/(\lambda - 1) < R < r_{i,1}$ ,  $i = 1, 2$ , then  $f_{\lambda,p_1} \neq f_{\lambda,p_2}$  by Remark 1.1, while  $\int_0^R f_{\lambda,p_i}(x) dx = \int_0^R u_{\lambda,p_i}(x) dx = 1$ ,  $i = 1, 2$ .

*Remark 2.2.* In the definition (1.1), let  $m = \infty$  with the condition  $\sum_{j=1}^{\infty} |c_{j+1} - c_j| < \infty$ . Then the derivative of  $p$  in the sense of distribution is a finite Radon measure and the Fourier transform of the derivative is an almost periodic function. With some conditions we can also construct solutions for (2.1) from such functions. For the Fourier preimage of the space of all finite Radon measures, see, for example, [6, 7].

In the rest of this section, we give graphs of solutions for  $f'(x) = 4f(2x)$ ,  $f'(x) = (3/2)^2 f(3x/2)$  and  $f'(x) = 9f(3x)$  with  $f(0) = 0$ .

Let  $p$  be as in Table 1. Then, calculating  $T_{\lambda,p}^\ell \chi_{[0,1]}$  numerically by computer, we get the graphs of solutions (S1)–(S7) in Figure 1, (S8)–(S11) in Figure 2 and (S12)–(S13) in Figure 3.

## REFERENCES

- [1] A. Augustynowicz, H. Leszczyński and W. Walter, *On some nonlinear ordinary differential equations with advanced arguments*, Nonlinear Anal., 53 (2003), 495–505.
- [2] V.I. Burenkov, *On estimates of Fourier transforms and convolutions in Nikol'skij-Besov spaces*, Proc. Steklov Inst. 3(1990), 35–44






FIGURE 1. Solutions of  $f'(x) = 4f(2x)$ .

$\lambda$	$p \in X$	Solution
$\lambda = 2$	$\chi_{[0,1)}$	(S1)
$\lambda = 2$	$\chi_{[0,1/2)} + (3/4)\chi_{[1/2,7/6)}$	(S2)
$\lambda = 2$	$\chi_{[0,1/2)} + (1/2)\chi_{[1/2,3/2)}$	(S3)
$\lambda = 2$	$\chi_{[0,1/2)} + (1/4)\chi_{[1/2,5/2)}$	(S4)
$\lambda = 2$	$\chi_{[0,1/2)} + (5/4)\chi_{[1/2,9/10)}$	(S5)
$\lambda = 2$	$\chi_{[0,1/2)} + (3/2)\chi_{[1/2,5/6)}$	(S6)
$\lambda = 2$	$\chi_{[0,1/2)} + 2\chi_{[1/2,3/4)}$	(S7)
$\lambda = 3/2$	$\chi_{[0,1)}$	(S8)
$\lambda = 3/2$	$\chi_{[0,1/2)} + (5/4)\chi_{[1/2,9/10)}$	(S9)
$\lambda = 3/2$	$\chi_{[0,1/2)} + (3/2)\chi_{[1/2,5/6)}$	(S10)
$\lambda = 3/2$	$\chi_{[0,1/2)} + 2\chi_{[1/2,3/4)}$	(S11)
$\lambda = 3$	$\chi_{[0,1)}$	(S12)
$\lambda = 3$	$\chi_{[0,1/2)} + (1/4)\chi_{[1/2,5/2)}$	(S13)

TABLE 1.  $\lambda$ ,  $p$  and the solution  $f_{p,\lambda}$ .FIGURE 2. Solutions of  $f'(x) = (3/2)^2 f(3x/2)$ .FIGURE 3. Solutions of  $f'(x) = 9f(3x)$ .

- [3] L. Fox and D. F. Mayers, *On a functional-differential equation*, J. Inst. Maths Applics, 8 (1971), 271–307.
- [4] P. O. Frederickson, Global solutions to certain nonlinear functional differential equations. J. Math. Anal. Appl. 33 (1971), 355–358.
- [5] P. O. Frederickson, Dirichlet series solutions for certain functional differential equations. Japan-United States Seminar on Ordinary Differential and Functional Equations (Kyoto, 1971), 249–254. Lecture Notes in Math., Vol. 243, Springer, Berlin, 1971.
- [6] Y. Giga, K. Inui, A. Mahalov and S. Matsui, Uniform local solvability for the Navier-Stokes equations with Coriolis force, Methods Appl. Anal. 12 (2005), 381–393.
- [7] Y. Giga, A. Mahalov, H. Jo and T. Yoneda, On time analyticity of the Navier-Stokes equations in a rotating frame with spatially almost periodic data, preprint.
- [8] M. Heard, *A family of solutions of the IVP for the equation  $x'(t) = ax(\lambda t)$ ,  $\lambda > 1$* , Aequationes Math., 9 (1973), 273–280.
- [9] A. F. Ivanov, Y. Kitamura, T. Kusano and V. N. Shevelo, *Oscillatory solutions of functional differential equations Generated by deviation of arguments of mixed type*, Hiroshima Math. J. 12 (1982), 645–655
- [10] T. Kato *Asymptotic behavior of solutions of the functional differential equation  $y'(x) = ay(\lambda x) + by(x)$* , in: Delay and Functional Differential Equations and Their Applications, Proc. Conf., Park City, Utah, 1972, Academic Press, New York, 1972, 197–217.
- [11] T. Kato and J. B. McLeod, *The functional-differential equation  $y'(x) = ay(\lambda x) + by(x)$* , Bull. Amer. Math. Soc. 77 (1971), 891–937.
- [12] T. Kusano, *Oscillation of even order Linear Functional Differential Equations with Deviating Arguments of Mixed Type*, J. Math. Anal. Appl. 98 (1984), 341–347.
- [13] E. Nakai and T. Yoneda *Construction of solutions for the functional-differential equation  $f'(x) = af(\lambda x)$ ,  $\lambda > 1$ , with  $f(0) = 0$* , preprint.
- [14] J. R. Ockendon and A. B. Tayler, *The dynamics of a current collection system for an electric locomotive*, Proc. Roy. Soc. Lond. A. 322, 447–468 (1971).
- [15] Y. Sawano and T. Yoneda, *On the Young theorem for amalgams and Besov spaces*, International Journal of Pure and Applied Mathematics., 36, No 2 (2007), 199–208.
- [16] Y. Sawano and T. Yoneda, *Quarkonial decomposition suitable for integration*, to appear in Mathematische Nachrichten.
- [17] H. Triebel, *Theory of function spaces*, Monographs in Mathematics, 78. Birkhser Verlag, Basel, (1983).
- [18] W. Volk, *Properties of subspaces generated by an infinity often differentiable function and its translates*, ZAMM Z. Angew. Math. Mech. 76 (Suppl. 1) (1996) 575–576.
- [19] W. Volk, *Evaluation of the solution of an integral-functional equation*, Journal of Computational and Applied Mathematics 174 (2005) 423–436.
- [20] T. Yoneda, *Spline functions and  $n$ -periodic points (Japanese)*, Transactions of the Japan Society for Industrial and Applied Mathematics (Japanese), 15 (2005), 245–252.
- [21] T. Yoneda, *On the functional-differential equation of advanced type  $f'(x) = af(2x)$  with  $f(0) = 0$* , J. Math. Anal. Appl., 317 (2006), 320–330.
- [22] T. Yoneda, *On the functional-differential equation of advanced type  $f'(x) = af(\lambda x)$ ,  $\lambda > 1$ , with  $f(0) = 0$* , J. Math. Anal. Appl., 332 (2007), 487–496.

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, THE UNIVERSITY OF TOKYO,  
3-8-1 KOMABA, MEGURO-KU TOKYO 153-8914, JAPAN  
*E-mail address:* yoneda@ms.u-tokyo.ac.jp